## **BUSD Illness Protocol**

## I need to stay home if:

	54		00	(65)		
Fever	Vomiting	Diarrhea	Rash	Head Lice	Eye	Hospital
					Infection	stay
						and/or ER visit
Temperature of 100 or higher	Within the past 24 hours	Within the past 24 hours	Body rash with fever	Itchy head, active lice	Redness, itching, and "crusty" drainage from eye	Hospital stay and/or ER visit
I am ready to come back to school when:						
Fever free for 24 hours	Free from	Free from	Free from	Treated with	Evaluated by	Released by medical
without the	vomiting for 24 hours.	diarrhea for at least 24	rash or fever. Have been	appropriate lice	my doctor and have	provider to
use of fever	24 110013.	hours.	evaluated by	treatment at	note to	return to
reducing			doctor if	home and	return to	school.
medication,			needed.	checked in	school.	
i.e. Tylenol or				the office		
Motrin.				upon return		
				to school.		

CDC always recommends everyday preventative actions to help prevent the spread of viruses including:

- Wash your hands often with soap and water for at least 20 seconds. If soap and water are not available, use an alcohol based hand sanitizer.
- Avoid touching your eyes, nose, and mouth with your unwashed hands.
- Avoid close contact with people who are sick.
- Stay home when you are sick.
- Cover your cough or sneeze with a tissue then throw the tissue in the trash.
- Clean and disinfect frequently touched objects and surfaces.

These are everyday habits that can help prevent the spread of viruses.